

# Conservation in Qatar:

## Impacts of Increasing Industrialization

---

*Renee Richer*



**CIRS**

CENTER FOR  
INTERNATIONAL  
AND REGIONAL  
STUDIES

---

GEORGETOWN UNIVERSITY  
SCHOOL OF FOREIGN SERVICE IN QATAR

2008

# Conservation in Qatar: Impacts of Increasing Industrialization

Renee Richer

Center for International and Regional Studies  
Georgetown University School of Foreign Service in Qatar

2008



# CIRS

CENTER FOR  
INTERNATIONAL  
AND REGIONAL  
STUDIES

Established in 2005, the Center for International and Regional Studies at the Georgetown University School of Foreign Service in Qatar is a premier research institute devoted to the academic study of regional and international issues through dialogue and exchange of ideas, research and scholarship, and engagement with national and international scholars, opinion makers, practitioners, and activists.

Guided by the principles of academic excellence, forward vision, and community engagement, the Center's mission revolves around five principal goals:

- To provide a forum for scholarship and research on international and regional affairs
- To encourage in-depth examination and exchange of ideas
- To foster thoughtful dialogue among students, scholars and practitioners of international affairs
- To facilitate the free flow of ideas and knowledge through publishing the products of its research, sponsoring conferences and seminars, and holding workshops designed to explore the complexities of the twenty-first century
- To engage in outreach activities with a wide range of local, regional and international partners.

This publication series is made possible by the generous support of Qatar Foundation for Education, Science and Human Development.

# Conservation in Qatar: Impacts of Increasing Industrialization

Renee Richer

*Renee Richer joined Weill Cornell Medical College in Qatar in January 2007, where she teaches biology. Richer received her B.A. in biology from the University of Chicago and her Ph.D. in biology from Harvard University in 2004. Her research experiences range from animal physiological ecology to plant physiological ecology. Richer's dissertation work focused on climate change and savanna ecosystems in southern Africa. She was previously an assistant professor and Director of the Environmental Conservation and Research Center at the American University of Armenia. Her work with bird life in Armenia was recognized in 2007 by the Whitley Award, the United Kingdom's largest conservation award.*

# Foreword

Like all industrial countries, Qatar is facing environmental degradation and the challenges posed by the loss of ecosystem services. However, unlike many countries, Qatar is extremely small, hot and arid, with few obvious natural resources other than abundant gas and oil reserves. It is almost completely dependent upon imports for food and other commodities to support the burgeoning population. Social and economic changes are taking place at an alarming rate, putting at risk the natural and cultural resources of Qatar. However, such loss of natural and cultural heritage need not be the case and great economic benefits can be gained from ecologically based development. Qatar is in a unique position, given the financial resources and forward thinking leadership, to move forward and be amongst the first countries ready to take advantage of the next economic revolution: the green revolution.

This review paper is a continuation of the Center for International and Regional Studies at the Georgetown University School of Foreign Service in Qatar's panel presentation on the environment. The Panel presenters addressed a variety of environmental issues facing in Qatar in light of increasing industrialization. The program "Environmental Degradation and Conservation: Challenges and Prospects" was held on November 12, 2007. The presentation and more importantly, the fact that attention to environmental issues is growing, was overwhelmingly well received by academics and the public.

In response to the demand for greater awareness of environmental issues and more environmentally related work regarding Qatar, this review paper was developed. The goal is to highlight some of the major environmental issues, the challenges and opportunities in Qatar, as well as increasing the general knowledge, awareness of and action on the environment. It is a collaborative effort between CIRS and Dr. Renee A. Richer of Weill Cornell Medical College in Qatar.

Renee Richer  
Weill Cornell Medical College in Qatar

Despite its meteoric rise in importance and its newly acquired urgency in global politics and academic studies, environmental scholarship is a largely understudied and overlooked subject in Qatar. It was this scholarly vacuum that prompted us to embark on a collaborative, multi-disciplinary endeavor to better understand the environmental challenges facing the Gulf region in general and Qatar in particular, an important part of which is presented here in the form of this Occasional Paper. I am thankful to Renee Richer for agreeing to take on this important undertaking.

This Occasional Paper fills a significant gap in the academic literature on the relationship between industrial growth and the environment in Qatar. The study highlights the dangers of increased environmental degradation due to growing industrial development and human activities. Richer points out that although Qatar is an arid desert, there is a surprising abundance of life-forms that have adapted to the harshness of the climate. Richer warns that because biodiversity in Qatar is under-researched, it is also under-conserved. This could result in a number of negative consequences, not the least of which is the possible extinction of a number of species that have yet to be fully studied by scientists.

Perhaps the most important insight to be drawn from Richer's study is the fact that conservation is actually an economically viable ethos that increases, rather than decreases, business potential. Richer argues that with the necessary political will, Qatar could make great strides in and act as a pioneer of environmental conservation as well as satisfy industry demands. Richer insists that Qatar could protect its biodiversity and have thriving industries by investing in research geared towards the benefits of sustainable development.

Drawing on the insights of members of the scientific community, Georgetown University's Center for International and Regional Study will continue to engage in research and examination of environmental and policy issues of contemporary importance. Through such efforts, we hope to contribute to a deepening understanding of one of the most critical issues facing our generation.

Mehran Kamrava  
Director, CIRS

# Abstract

Industrial development in the State of Qatar is taking place at an unprecedented rate. Such development is putting the environment at the risk, threatening ecosystem services and biological diversity. While Qatar is currently developing the legislation, regulatory bodies and management agencies for successful ecosystem management and conservation efforts, the full implementation of these protective measures has yet to be achieved. This is in part due to a lack of scientific expertise and trained manpower as well as the early stage of their development. While strides have been taken by the State of Qatar, the question remains whether they can be implemented in a timely fashion to ensure that current and future development projects support the country's goal of sustainable development.

# Acknowledgements

Dr. Iain Macdonald contributed significantly to the preparation of this paper and the initial panel presentation on biodiversity and conservation in Qatar. His help is greatly appreciated. Dr. Benno Boer (UNESCO), S. Faizi (SCENR), Dr. Mohamed Alaa Abdel-Moati (SCENR) and Dr. Elsadig Bashir (FEC) supplied important resources and information, greatly enhancing the paper.

# Introduction

Biodiversity and Conservation are not necessarily the first things that come to mind when one visits Qatar. The stark landscape, punctuated by massive industrial sites, pipelines and refineries belies the diversity of organisms that Qatar supports and more importantly, the diversity that Qatar could support. That stark and seemingly barren landscape is not completely natural. It is partly the result of human activity and the activities of animals that are associated with humans. Unless measures are taken to halt further degradation and maintain the current conditions or preferably, restore the environment, there is likely to be irrevocable environmental degradation. Qatar is not unique in the environmental challenges it faces, but it is unique in the role that it plays in providing habitats for species vulnerable and threatened with extinction and the species yet unknown.

The Arab world, in general, has identified the environmental challenges of water scarcity and desertification or land degradation as the most pressing environmental issues in the region. In fact, the challenges go beyond these two issues (El-Sayed Selim, 2004). Like all countries, Qatar faces threats from invasive species, climate change, pollution and habitat change, the four main threats to global biodiversity identified in the Convention of Biological Diversity. Most of these threats are a result of industrialization and the changes that accompany industrialization from within and outside the country.

## *What Makes Qatar Unique?*

The State of Qatar is a small peninsula, just over 11,000 km<sup>2</sup> with 900 km of coastline, off the Arabian peninsula and connected in the south to Saudi Arabia. It is generally flat and rocky, with sand dunes in the south of the country. Qatar is hot, dry and surrounded by a shallow, hyper-saline, semi-enclosed sea. As a subtropical desert, the annual rainfall is a mere 81mm with an average annual maximum temperature of 31°C, although absolute maximum air temperatures can exceed 47°C. The high temperature, strong winds, low rainfall and low nutrient availability of the soil mean that recovery of the terrestrial ecosystems from disturbance is very slow. This makes the Arabian peninsula and Qatar in particular one of the most hostile environments on earth (Brook et. al. 2006), and one of the most fragile.

High salinity and great temperature fluctuations make the Gulf surrounding Qatar unique and extreme. Tremendous solar pumping (evaporation) and the lack of water exchange with the open ocean makes the Gulf saline and hot, and the lack of exchange means that pollutants in the Gulf take longer to dissipate (Brook et. al. 2006). The local marine resources are typically surviving at their extreme tolerances of environmental parameters; however, this does not stop them from displaying the ability to recover from mass mortality events (Shinn 1976). The Gulf is relatively small and shallow with an average depth of only 35 meters and the greatest depth is only 120 meters. On the west coast, between Qatar and Bahrain, the average water depth ranges from 1-5 meters. There is little water exchange with the northern Indian Ocean through the Strait



of Hormuz. Therefore, the Gulf has the largest annual temperature variation of any sea supporting corals. Water temperatures vary between below 11-15 °C in winter and can reach as high as 34-40 °C in summer (Rezai et. al. 2004). While the salinity levels of the open ocean are approximately 35 ppt (parts per thousand), those of the Gulf are around 45 ppt and 70 ppt in the shallow areas off the northwest coast of Qatar. In the Gulf of Salwa, on the west coast of Qatar, salinity and temperature extremes may be even higher. Salinity levels as much as 200 ppt have been recorded in other areas of the Arabian Gulf (Carpenter et. al., 1997).

Qatar is home to 1955 known species; 955 are marine. These species are specially adapted to the unique hyper-arid, hypersaline, and thermovolatile conditions. While these extreme environmental factors were previously believed to limit biodiversity in the region, the number of species recorded in Qatar continues to increase (Macdonald, 2007), some of those being new to science. Ironically, this extreme habitat is likely to be home to species with unique adaptations to the extreme conditions and a rich genetic resource for biotechnological advances. The most thermally tolerant coral species, i.e., those that don't bleach when other species do in the Gulf, are found here and they host a large proportion of symbiotic zooxanthellae compared to other corals around the region. Interestingly, some species of endemic bacteria in the Gulf can utilize hydrocarbons (oil) (Radwan et. al., 2005). This is thought to be a result of the natural oil seeps in the Gulf region, resulting in a relatively high level of hydrocarbons in the water providing a selective pressure for utilization or sequestration of the hydrocarbons. Such biological resources could ultimately be used to develop oil spill remediation measures.

Biodiversity is often grossly undervalued by society and the institutions responsible for making decisions that threaten biodiversity (Dietz and Adger, 2003). This is in large part attributed to the lack of knowledge and understanding of the benefits that biodiversity provides in terms of ecosystem services. This lack of knowledge, as well as knowledge of the species and habitats, limits conservation efforts in Qatar. While some forms of environmental degradation can be remediated, biodiversity loss is considered irreversible (Dietz and Adger, 2003).

## *The Challenges of Industrialization*

Increasing industrialization poses many different challenges to the environment. These are often associated with large-scale energy production and metallurgy but also include the social and economic changes that accompany technological innovation. While there are the obvious concerns about emissions, waste management and land degradation associated with industry, one important aspect to consider, particularly in Qatar, is the increase in population that comes with increasing industrialization. An increasing population places demands on land for housing and recreation, waste disposal and sewage treatment and other resources such as infrastructural improvements. (Yasseen and Al-Thani, 2007).

Secondly, development has occurred at an unprecedented pace. In the matter of a few years Qatar has gone from being one of the poorest countries to being one of the richest countries in the world. Because this development has taken place at such a fast rate, there is simply a lack of capacity in regulatory and monitoring bodies to help guide the growth. Only in 1986 was an Environmental Protection Committee established which was replaced by the Supreme Council for the Environment and Natural Resources (SCENR) in 2000. In that same year a draft environmental law was estab-

lished with 100 articles, making environmental policy and implementation relatively new to Qatar.

Until relatively recently, Qatar was dominated by nomadic and semi-nomadic people whose livelihood depended on fishing, pearling, camel breeding and dhow construction. However, the discovery of oil has spurred not only socio-economic changes, but also environmental changes (Khan, 2002). The tremendous economic boom seen in Qatar has been supported largely by an influx of expatriate workers. With the fastest growing economy in the Arab world and the world's third largest gas reserves, Qatar's GDP has been growing at an annual rate of 18% between 1999 and 2004. However, between 2003 and 2004 the annual increase in GDP was 21% with the 2005 GDP over \$47,000 US dollars (Planning Council, 2007). Crude oil and gas account for 62% of the Qatari GDP. In the oil and energy sector alone, the Qatari population is not large enough to support the labor demand and it is unlikely they will be able to meet that demand in the future. The annual population growth for non-Qataris is 6.5% while that for Qataris has been 2.2% between 1997 and 2004 (World Bank, 2005). In sharp contrast, 40% of expatriates in Qatar fell below the poverty line compared with only 1% of Qataris (Khan, 2002). Such poverty among the workforce may exacerbate environmental pressures.

The annual increase of expatriate workers is likely to have increased significantly since this last census. In fact, World Bank (2005) estimates of population growth for Qatar were to reach approximately 600,000 by 2006. In actuality, the population increased to nearly 800,000 by that time. Predictions by the UN for Qatar for the year 2025 placed a high estimate at 850,000 while that of the Planning Council for 2015 was 1.4 million (Planning Council, 2007). However, both predictions were gross underestimates because by the end of 2007 the population had reached 1.5 million (Gulf Times, 2008).

## Qatar's Natural Resources

### *Marine Resources*

Qatar's marine environment supports 955 known marine species. These include 3 documented mammals (although more than 20 are known in the region), 15 reptiles, 136 fish species and 379 species of plants. The marine environment includes coral communities, intertidal mud flats, algal beds, seagrass beds, pearl oyster beds and mangroves. These habitats, generally the most productive, are located in waters from the shoreline to only 10-12 meters deep, and therefore closest to areas of human activities (Khan, 2007). They support a range of marine mammals including dugong, and about twenty species of whales and dolphins including the Humpback whale, Brydes whale and the Bottlenose dolphin. Sharks are common and include the Tiger shark, Hammerhead shark, and the White shark (SOMER, 2003).

Eighty percent of the mangroves in the Gulf occur on the Iranian side of the Gulf. However, mangrove (*Avicennia marina*) occurs along the coast of Bahrain, Qatar and the United Arab Emirates while not generally found in Kuwait or Saudi Arabia (Al Habshi et. al., 2007). The mangrove stands stabilize the sediments, protect the coastline, provide shelter for animals, particularly commercially important juvenile fish, and their leaf fall provides an important food source for marine species (Brook et. al.,

2007). Mangrove habitats in general are globally threatened as a result of coastal development, fish and shrimp farms and other environmental stressors.

The seagrass beds, the most productive of the marine habitats especially for commercially important marine species, occur in shallow, protected waters. They support a wide variety of organisms, over 600 animal species recorded (SOMER, 2003), including molluscs, crustaceans, fish, turtles and the unique marine mammal, the dugong (*Dugong dugon*) (Loughland et. al., 2007, Brook et. al., 2006). Known as the sea cow, the dugong, feeds by grazing on the seagrass beds in the shallows. The dugong is the only species within the Gulf to be highlighted as an EDGE species (Evolutionary Distinct and Globally Endangered) a scheme set up by The Zoological Society of London. At one time the dugong was actively hunted for food. That has now decreased with industrialization (i.e., the lack of need for subsistence fishing) and the introduction of local regulations to prevent such hunting. However, dugongs are still accidentally caught in nets, hit by small boat propellers and may be hunted clandestinely (Sheppard et. al. 1992). A large herd of over 600 individuals has been observed off the west coast of Qatar in the Gulf of Salwa. Estimated at 7,000 individuals, the Gulf population is the world's second largest population of dugongs and is important for the continued success of the vulnerable species (SOMER, 2003). However, with the long gestation and calf period, population replenishment is slow. A Memorandum of Understanding (MoU) signed by the UAE and other States such as Australia, France and Iran (November 1, 2007) was designed to protect the dugong and its habitat. Unfortunately, signatories of the MoU failed to include Bahrain and Qatar (Khaleej, 2007).

Five species of sea turtle are known in the Gulf. These include the globally endangered Green turtle (*Chelonia mydas*), critically endangered Hawksbill turtle (*Eretmochelys imbricata*), critically endangered Leatherback turtle (*Dermochelys coriacea*), endangered Loggerhead turtle (*Caretta caretta*) and endangered Olive Ridley turtle (*Lepidochelys olivacea*) (SOMER, 2003). The Green Turtle feeds within the seagrass beds and the Leatherback Turtle has been observed on the coast of Qatar but is not officially recorded. The Hawksbill Turtle is also known in Qatar and is reported to breed on various offshore islands (SOMER, 2003) and along the northeastern coastline on suitable beaches. In general, the males tend to stay offshore and the females come to shore for nest production and egg laying. Unfortunately, despite protective measures, turtle eggs are still collected along the shore.

The shallow water surrounding Qatar supports a large coral community. The corals may extend as much as 48-80 km from the shoreline (Ahmed, 2002). Because the water is shallow, light can penetrate to support photosynthetic organisms. This light ultimately provides the energy to support the entire community. However, only eight species of coral have been reported for Qatar. This seems unusual given that 31 species are recorded in Bahrain and 34 species in the United Arab Emirates (SOMER, 2003). This low species count could be attributed to lack of adequate data in the region or natural causes such as salinity.

Finally, marine species are important commercially. Species such as lobster, octopus and many local fish species such as grouper and bream are locally exploited (Ahmed, 2002). Landings of emperor (*Lethrinus* sp.) and grouper (*Epinephelus* sp.) accounted for 42% of total landings in 2001 (FAO, 2003). Shrimp exploitation has been stopped in order to increase the local population. However, given that Bahrain and Saudi Arabia still catch shrimp, recovery of the population is unlikely under these circumstances. Qatar manages the fishing industry by regulating the inputs, i.e. the

number of artisanal vessels, rather than limiting the catch size. In 1995 the number of vessels was 493 and between 2001 and 2003 the number of fishing vessels stayed constant at 515. However, small vessels could be replaced with larger vessels and 49% of the vessels are 16.4 m greater in length (FAO, 2003). This may be responsible for the rise in the number of fishermen increasing from 3,101 in 1995 to 4,721 in 2001. The total annual fish catch also increased from approximately 70,000 tons to 140,000 tons between 2000 and 2005, an annual increase of over 14%. It is important to note that Qatar has no seasonal ban on fishing (Peninsula, 2007).

## *Terrestrial Resources*

Qatar's terrestrial environment supports 1,000 known species. These include 8 mammals, 255 birds, 29 reptiles, one amphibian and 228 invertebrates. There are 317 species of plants from 61 families and 142 known species of fungi. Of the terrestrial species about 26 are at risk. These include the Greater spotted eagle, Lesser kestrel, Socotra cormorant and Houbara bustard. Arabia's only antelope, the Arabian oryx (*Oryx leucoryx*), inhabited the deserts until a century ago (Brook et. al. 2006). It is now extinct in the wild but re-introduction programs are attempting to introduce it, along with the Reem gazelle (*Gazella leptoceros*), back into the wild (Brook et. al., 2006). Other mammals present in Qatar include the Desert hare (*Lepus capensis*), Ethiopian hedgehog (*Paraechinus aethiopicus*), and two species of fox: the Arabian fox (*Vulpes vulpes Arabica*) and the Sand fox (*Vulpes ruppellis*).

Bird surveys, which unfortunately are dated, suggest there are approximately 255 bird species recorded in Qatar. Of those, 23 are local breeders, 78 are winter visitors and 50 are vagrants. The remaining 104 species, 44% of the species known in Qatar, are migrants which simply pass through Qatar (FEC). Updated information should be available soon, pending a recent survey supported by industry. The invertebrates, which comprise the majority of the animal life, are little studied and comprehensive lists of species or groups are not available. In terms of terrestrial plants, seven species of plant are present in Qatar that are known to be threatened with extinction (Planning council, 2006).

## **Environmental Risks in Qatar**

Environmental risks range from the global risks associated with large-scale international actions to regional actions and ultimately to local activities that can be modified or controlled. Although Qatar currently has the largest CO<sub>2</sub> per capita emission rate at 69 tons CO<sub>2</sub>/per person/per year, which dwarfs the second largest emission rate of 39 tons CO<sub>2</sub>/per person/per year by the UAE (WRI, 2007), that number reflects the fact that Qatar produces a vast amount of the world's energy resources and has a relatively small population. However, in May of 2007 Qatar Petroleum successfully registered the first United Nations Clean Development Mechanism (CDM) project under the Kyoto protocol in the Gulf region (CDM, 2007). That project is designed to decrease CO<sub>2</sub> emissions by reducing flaring.

Globally, current climate change predictions for the Gulf region vary based on different models and the IPCC scenario, especially for precipitation (Chaudhary and Husain, 2006). Because precipitation is influenced by local conditions and complicated teleconnections, predictions for precipitation changes are particularly difficult to construct.

Model predictions for temperature and humidity all show an increase for the Arabian peninsula, the only question being how large of a temperature increase (Chaudhary and Husain, 2006). Since Qatar is a desert region with low annual precipitation, small changes in precipitation, either an increase or decrease, can be expected to cause significant changes in species associations for both the marine and terrestrial environments. As disease vectors are also expected to change, the incidence and occurrence of disease may play a significant role in altering species associations (Chaudhary and Husain, 2006).

### *“Water, water everywhere, nor any drop to drink”*

One of the most critical issues associated with increasing industrialization and the concomitant population increase is the availability of water. Qatar is well below the World Bank “water poverty line” of 1000 cubic meters per person per year, with only an average of 91 cubic meters available per person per year (El-Sayed Selim, 2004). On the Arabian peninsula in general, fossil water resources are believed to have accumulated during the pluvial period from 10,000-30,000 years ago (Alsharhan et. al., 2001). Current rate of water use is about six times the natural renewal rate of water resources while over half of the water used is desalinated water or treated wastewater (Brook, Houqani and Mugrin, 2006). Nearly 96% of the domestic water in Qatar is provided by desalination, while ground water supplies are largely used for the small agricultural industry. Continued over-exploitation of the fossil water reserves threatens the remaining reserves from saltwater intrusion while over-use of ground water for agriculture is resulting in soil salinization and land degradation (Child, 2006).

Over-exploitation has resulted in increasing salinity levels in ground water, from saltwater intrusion, from 71-1160‰ (Brook et. al., 2006). Importantly, over-use of ground water has resulted in the disappearance of many springs in the Arabian peninsula in general. These natural oases were an important habitat for reptiles, birds and mammals including rare and endemic species. As demands increase, the need for desalinated water will increase. Demand simply cannot be met by renewable or fossil resources. Currently among the GCC countries there are 15 desalination plants located on the Gulf, releasing gases, hot brine, treatment chemicals and other trace elements (Abderrahman and Husain, 2006). At this time, the effects of the desalination plants on the Gulf as a whole are limited in scope, but the effects are greater locally. As the number of desalination plants increases, their impact on the sensitive marine environment may become devastating. A recent study in Kuwait, showed that the hot water released by a power and desalination plant reduced sea grasses and coral reefs by 15-20% just from the increase in temperature alone (Abderrahman and Husain, 2006).

### *Threats to the Marine Environment*

The challenges to the marine environment also include global changes such as climate change, but also local effects of industrialization from land-based sources such as the releases from power and desalination plants and sewage treatment plants, pollution from the petrochemical industry, coastal reclamation projects, ballast water, invasive species, eutrophication, overfishing and threats from increased tourism (Khan, 2007).

Coastal development is a major threat to the marine environment. Coastal development such as dredging and land reclamation for projects such as residential areas, industry, causeways, fishing ports, airports and harbours takes its toll (SOMER, 2003). Dredging

has been important for providing access to major ports for shipping. However, these activities result in siltation and increased water turbidity, which can inhibit photosynthesis, clog fish gills and essentially smother the algal mats, seagrasses and coral communities (Al-Ghadban and Price, 2002). This is a continuing stressor on the coral communities and the intertidal and shallow tidal ecosystems. In the case of corals, the increase of sediment in the water column decreases light levels and otherwise affects the health of the coral and can also prevent coral recovery (Rezai et. al. 2004). Coastal development utilizing infill and dredging, such as the artificial island developments, affects the shores. This has stressed fishing stocks as a result of damage to nursery and feeding grounds of the fish (Jones et. al., 2007). The causeway, currently under construction from Bahrain to Qatar, will utilize reclaimed areas of coral from Fasht Al Adhom. Besides the obvious damage to the coral community, water currents may also be disrupted in the Gulf of Salwah (Rezai et. al., 2004). Much of the development taking place on the coastal areas of Doha lacks appropriate impact assessments or restoration plans (Yasseen and Al-Thani, 2007). This development affects mangroves (*Avicennia marina*) and as many as sixteen species of xerophyte and halophytes.

Because the water surrounding Qatar is shallow with little exchange, characteristic of the Gulf in general, this area is susceptible to temperature extremes. Coral bleaching events, as a result of high sea temperatures, occurred in 1996, 1998 and 2002, reducing live coral cover to as little as 1% in shallow areas. The temperature anomaly in 1998, when temperatures exceeded 37.3°C in central regions of the Gulf, was more than 2°C above average (Riegl, 2002). This was the largest temperature rise in the southern Gulf since 1870 and emphasizes the increase in sea surface temperature in the Gulf of at least 0.2 °C per decade for the last 50 years (Sheppard and Loughland, 2002). Recovery has been largely limited to deeper water areas with less human impact. Estimates of reef destruction are as low as 1% for Oman and as high as 97% for Bahrain. Along with the loss of coral was a significant decrease in fish stocks and species richness (Riegl, 2002).

Although the awareness of the importance of coral conservation is increasing in the region, conservation in these areas still lags behind much of the world (Rezai et. al., 2004). As climate change continues and temperatures increase, it is questionable whether the corals will be able to recover. Most likely there will be changes in species composition and cover (Sheppard, 2003). Since 1950 extreme weather events, both hot and cold, have been increasing significantly in the Middle East region (Zhange et. al., 2005). Qatar experienced an increase of 0.10°C in the 20th century and the predicted increase in the 21st century ranges from 3.0-5.9 °C (Mitchell and Hulme, 2000). Industrial cooling water is also a threat to the marine environment. Cooling water is often discharged into the sea from co-generation desalination and power plants. The water can reach 10°C above ambient and contains chlorine and other chemicals to prevent scaling. In 2006 the “International Cooling Seawater Specialists and Operators Conference” was held in Doha to discuss temperature and chlorination challenges posed by cooling water. With a focus on Qatari industries, local experts developed management recommendations for SCENR (Peninsula, 2006). Important to note is that recent adoption of new water treatment techniques (i.e. pulse chlorination) has proven successful in decreasing, by as much as 56%, the amount of chlorine in cooling water releases. Driven by increasingly strict regulation concerning chlorine release, QatarGas has recently implemented this BAT (Best Available Technology) by the European Union (QatarGas, 2007). Other threats to the coral communities of Qatar include physical damage from anchors, propellers, abandoned fishing nets, traps and overzealous sport divers (SOMER, 2003).

Finally, as the demand for gas and oil increases, the transport of both products will increase through the Gulf. Currently over 60% of the world's total crude oil exports are transported through the Strait of Hormuz. Oil tankers pollute in many ways, including exhaust, routine leaks, ballast exchange, and spills, both war and error related. If we combine these risks with pipeline ruptures and well blowouts, this makes the Strait one of the world's highest oil pollution risk regions (SOMER, 2003). Given that the water exchange in the Gulfs is so slow and pollutants are slower to dissipate, spills can be devastating to the Gulf environment. The annual spill rate into the Gulf is 1.2 million barrels of oil from marine transportation (SOMER, 2003). Even now, more than fifteen years after the Gulf war, when more than 9 million barrels of oil were spilled in the Gulf, the Gulf environment has yet to completely recover from this disastrous oil spill (Barth, 2007). A comprehensive study of the effects of that spill have not been done in Qatar, and restitution has not been claimed by Qatar.

Ballast water from tankers poses a threat in the release of invasive species. As ships move globally from region to region, they often release non-native organisms into the water, disrupting local communities as well as introducing new species that can pose expensive problems for industry (Hushak and Deng, 1997).

As the population increases, discharge into the sea will likewise increase. In the case of sewage, approximately 20-30% of sewage discharged into the Gulf is not treated or only partially treated, promoting eutrophication (SOMER, 2003). At one time Qatar disposed of the domestic sewage directly into the Gulf. However, as of 1999, Qatar had 13 sewage treatment plants and claimed to no longer be dumping sewage. In addition, the treated water from the sewage treatment plants is used for irrigation of fodder crops, gardens, and city landscaping (SOMER, 2003).

## *Desertification in the Desert?*

The State of Qatar has made moves toward protecting its natural environment. These include public awareness campaigns, implementing a legal structure for the protection of species and habitats and by participating in regional programs like ROPME and the Convention on Biological Diversity. However, large-scale land reclamation projects, dredging and the discharge of cooling water will remain important issues for the marine environment (Rezai et. al. 2004).

It may seem ironic that Qatar, party to the Convention to Combat Desertification, is itself suffering losses due to desertification. How can desertification happen in the desert? It is important to understand the process of desertification and the impacts associated with desertification. The United Nations Convention to Combat Desertification defines desertification as land degradation in arid, semi-arid and dry sub-humid regions, partly as a result of fluctuating climate but also as a result of human agricultural activities such as grazing and crop production. Land degradation can include the loss of plant productivity, biodiversity and soil fertility.


Much of the Arabian Peninsula, including Qatar, has suffered from livestock grazing well above the ecological carrying capacity. To date there are 925 productive farms in Qatar, located on 9,800 hectares of land (Brook et. al., 2006). Although agriculture accounts for only 1% of GDP, agriculture accounts for 74% of freshwater use (Al Yousef et. al., 2000). In the past, the number of animals a family could support depended on their ability to provide water. Traditionally that involved labor-intensive maintenance of wells that often provided brackish water. Now that traditional herding

practices have changed combined with the collapse of the *hima* conservation system, there has been a decline of natural rangelands (Breulmann et. al., 2007). Today, fresh water can be easily transported and deep wells utilizing fossil ground water supplies are used to grow non-native feed and supply water which has allowed the livestock populations to increase dramatically. In addition, there has been a move away from nomadic life to a more sedentary life resulting in increased localized grazing pressure (Brown et al., 2006). Starting in the late 1950s the number of farms began to increase steadily such that by the late 1960s the effects of ground water over-exploitation was becoming evident (Al Yousef et. al., 2000).

Beyond water use, animals can place a tremendous stress on the local plant resources. The grazing and browsing pressure seen in Qatar is a result of not only camels but sheep and goats which, unlike camels, eat the plants down to the roots. As of 1999, Qatar supported nearly 4,000 horses, 14,000 cattle, 205,000 sheep, 175,000 goats and 47,000 camels (Al Yousef et. al., 2000) (Bourn, 2003). Qatar has the highest density of camels in the Middle East and the third highest amongst seventeen surveyed North African countries. The United Arab Emirates has the second highest camel density in the Middle East with nearly one half the density as Qatar (Brown et. al., 2006). With increasing industrialization, those animals will be further constricted on even less land, exacerbating what is already a significant problem. Such intensive grazing and browsing has resulted in the reduction of many plant species, changing the shrubland to the current, relatively barren, landscape witnessed in Qatar. Therefore, the population of Reem gazelles reintroduced into the Al Reem reserve must be supported with additional food resources (Brown et. al., 2006).

Before the grazing and browsing pressure increased, the plant species composition and habitat structure was significantly different with greater plant cover. Large areas of grassland once covered Qatar. Due to over-grazing, the grass community has largely been replaced by perennial shrubs that are thorny, toxic and generally unpalatable. The change from traditional to modern rangeland management has resulted in a decrease in vegetation density from 10% of land cover to approximately 1% of land cover. The change in cover associated with grazing and browsing animals can be illustrated by a satellite image of the Ras Laffan industrial area. The Ras Laffan industrial area is surrounded by a security fence, which also prevents the introduction of domesticated animals such as sheep, goats and camels. The satellite photo in Figure 1 illustrates the higher vegetation cover, with its darker green color, within the triangle of the security fence. It is likely that other secured areas may provide protection for vulnerable species. This is an important aspect of conservation that could be pursued.

Figure 1. Satellite image of area within security fence



Source: NASA, 2008 Terrametrics, DigitalGlobe, and Europa Technologies



This loss of vegetation cover and edible plant species as a result of grazing and browsing by camels and other domesticated animals has contributed to the disappearance of oryx and gazelles. The shrubland habitat at one time was able to support hyena, leopard, oryx and gazelle. Likewise, the loss of plant cover has led to an increase in wind erosion and a decrease in soil fertility, and all of these lead to declines in species diversity. Reintroduction of these species will require restoration of the shrubland habitat. This has been exacerbated by the fact that these large populations of domesticated animals are now supported by non-native species such as alfalfa and Rhodes grass which demand fresh water irrigation that promotes desertification (Breulmann et. al., 2007).

Restoration of the desert rangeland is possible in areas where the land has not been irrevocably damaged. This will require a commitment from all stakeholders as well as laws to manage grazing and the resources for full implementation of the program (Breulmann et. al., 2007). UNESCO is currently supporting a demonstration “Camel Farm” as a first step in the restoration of rangelands. Camels make ideal animals for the desert with their ability to drink water with a salt content up to 3% and eat native halophytes that would be toxic to other species. Unlike cows, they do not require air conditioning or sprinkling to keep cool, or large amounts of fresh water for drinking and fodder and they provide excellent meat and milk. In this way local fodder species could be produced that are less reliant on fresh water and pesticides and herbicides (Breulmann et. al., 2007). Camel farms would lead to a reduction in the number of camels on rangelands, allowing the habitat to recover.

## Ecosystem Management

Economic development need not be in conflict with environmental protection. In most cases we already have the technology available to be much more efficient. What is often lacking is the political willpower to lead. In fact, as we become increasingly aware of the true dollar value of ecosystem services and develop novel technologies and management techniques, environmentally friendly development is becoming much more economical. Full implementation ultimately involves the cooperation of industry, the public and government.

Successful conservation programs work with the local community, with regards to their values and practices, to protect the natural environment. In the case of Qatar, and Arabia in general, the unique concept of *hima*, an area set aside specifically for the sustainable use of natural resources by and for local communities, can be encouraged (Seddon and Mallon, 2007). How the traditional institution of *hima* can be used in the region is increasingly being debated among local and trans-boundary environmental protection bodies.

Coastal management plans have been developing in the region with the aid and support of stakeholders including NGOs, industries, and the general public. However, most of the coastal management plans have not been fully developed and full implementation has yet to be achieved (Krupp et. al., 2006). In addition, the use of Environmental Impact Assessments has been increasing. Environmental Impact Assessments is an evaluation of the impacts of a project on the environment, whether bad or good. The process includes not only an assessment of the effects on the environment but also the social and cultural impacts as well as suggestions or requirements for mitigation or bioremediation. In this way, decision-makers are fully informed of the expected im-

pacts of a project. Environmental Impact Assessments may ultimately result in alteration of the project, relocation or even abandonment of a poorly designed project. This is crucial given the coastal dredging and landfill projects including those currently taking place in Qatar largely associated with industrial, tourism and port developments.

It has been suggested that coastal development projects such as waterways, canals and islands can be designed to improve productivity. This could include spawning grounds and nursery areas (Krupp et. al., 2006). An example of this type of development is the West Bay Lagoon area in Doha. This development project introduced a new waterway, through the relatively unproductive *sabkha* creating intertidal and subtidal areas. The hope is that if carefully planned and managed, these waterways can act as a nursery for commercial fish species since seagrass species such as *Halodule uninervis* have been able to occupy these areas (Jones et. al., 2007). Although the seagrass species may be established, it is still unknown whether they will develop beds large enough to support other species.

## Environmental Impact Assessment and Industry

The Environmental Impact Assessment (EIA) also referred to as the Environmental, Social and Health Impact Assessment (ESHIA) process is a very important aspect of sustainable development and managed development. The Environmental Impact Assessment process in Qatar is under the review of the Supreme Council for Environment and Natural Resources (SCENR). Unfortunately, an official translation of the guidelines and criteria for EIA from Arabic to English, is currently not available in Qatar. Environmental Impact Assessments are overseen by the technical affairs section of the SCENR which is responsible for permitting. Initially, a Scope of Work is submitted to SCENR and then a decision is made as to whether or not the project requires a full EIA. If a project results in environmental damage, that damage must be remediated by the developer. In the event that the damage cannot be remediated, then the developer is responsible for providing compensation for the damage. Such compensation is not a fixed amount, rather it is a percentage of the project budget. The percentage is decided on a case by case basis. Ultimately the EIA committee, a team of five internal and external experts, makes a recommendation and SCENR has the final decision-making ability. However, there doesn't appear to be exact criteria stipulating which factors make a project successful or unsuccessful. That seems to be decided on a case by case basis depending on the scale and scope of the project.

However, given the pace of development and the need for EIAs, there is a demand for increased capacity in SCENR. More than 1,000 projects were submitted to SCENR for EIA in both 2006 and 2007 including large scale and medium/small scale projects. In many cases the company is asked to change the design or location of the project. In 2007, 5% of the projects did not pass the EIA. Currently the process is considered slow, taking one to two months to complete and resulting in delays. Therefore, SCENR stressed the great need for an increase in local experts (SCENR, pers. comm.) Although it is important to note that comparable EIA's for large scale industrial projects in other countries can take much longer to complete with final project approval taking years.

The Supreme Council for Environment and Natural Resources employs a surveillance department to detect any illegal projects and activities. Of the 650 Council employees, 300 are involved in the twenty-four hour a day surveillance department with 120 designated to marine activities and 180 designated for terrestrial activities. The

most common types of violations include use of illegal fishing nets and fishing in restricted areas, with two to three violations happening daily. Other common violations include illegal dumping, un-permitted jetties and dust and noise production at construction sites. Ultimately, the prosecution of environmental violations take place under the district attorney designated specifically for environmental affairs (SCENR, pers. comm.).

The mitigation costs assigned to industry often involve funding research projects, and conservation programs but may not result in remediation of the environment itself. These studies are generally not available to the public and are reserved for government and industry use. However, members of the public may request a study. In such cases a summary report may be issued to the inquiring person (FEC, pers.comm.). For example, Dolphin Energy developed a pipeline project from Ras Laffan. Remediation for that project involved funding a three-year project studying the coral communities from Qatar to UAE, including full mapping of the reefs. Similarly the Ras Laffan port expansion project will affect the seagrass beds and coral communities in the region. Compensation for that project will amount to \$50 million over a five-year period designated for conservation projects (SCENR, pers. comm.). Artificial reefs (Dolphin Energy and others) as well as coral translocation (Qatargas) are being pursued by industry to promote coral conservation and development, although the true impact of these measures are unknown.

Ras Laffan Environment Association (RLEA), a consortium of oil and gas companies, sponsors a variety of environmentally related work. RLEA is currently sponsoring the Qatar Bird Project. Funding for the first year of the project began in July 2007, and the first report was issued in December 2007. The goal of the project is to assess bird life and Important Bird Areas in Qatar. In 2007, Qatar became a member of Birdlife International, a globally recognized bird conservation organization. Friends of the Environment Center (FEC) are the local representatives. Birdlife International has developed the Important Bird Area (IBA) concept as a conservation tool. Each country may have designated IBAs and those areas will be the focus of conservation efforts in order to maximize the conservation of species. Based on dated surveys from the 1970s, 1980s, and 1990s, Birdlife International suggested five IBA sites in Qatar. The Qatar Bird Project is designed to review and reassess the status of the suggested sites and investigate potential new sites.

As of November 2007, FEC, with the aid of representatives of Birdlife International, completed an extensive survey confirming three of the five suggested IBAs as indeed being prime bird conservation sites in Qatar. Those sites include Al Thakira mangroves, Khor Al Udaid and Alshat Island. The two remaining sites have yet to be surveyed. In addition, five additional sites were recommended for IBA consideration. These include Um Tais, Ras Rukon, Abu Nakhla pond, Ashat Island and Shahanniya.

## National Biodiversity Strategy and Action Plan

The increasing international understanding and acceptance of the importance of biodiversity and ecosystem services for continued development has resulted in increased efforts to protect habitats and reduce biodiversity loss. Qatar, as part of this movement, is party to the Convention on Biological Diversity, Convention to Combat Desertification (CCD), select annexes of the International Convention for the Prevention of Pollution From Ships (MARPOL) and Convention on International Trade in Endangered Species

of Wild Fauna and Flora (CITES). Qatar will be hosting the CITES international conference in 2010, drawing attention to Qatar's unique environment. Most recently, Qatar ratified the Cartagena Biosafety protocol of the Convention on Biological Diversity in early 2007 and is now in the process of implementing the treaty.

The Convention on Biological Diversity (CBD) entered into force on December 29, 1993, with the goal of conserving biological diversity, sustainable use of species and habitats and the sharing of benefits derived from them. As party to the Convention on Biological Diversity, the State of Qatar has a legally binding obligation to implement the provisions of the Convention. Most signatory countries, of which Qatar is no exception, have Biodiversity Action Plans that outline the philosophy, policies and procedures of the country to successfully implement the Convention. The National Biodiversity Strategy and Action Plan (NBSAP) was completed in October of 2004 with the goal of reaching the 2010 Target, reducing the rate of biodiversity loss.

The NBSAP for the State of Qatar has eleven strategic goals encompassing the following themes:

- protected areas
- ecotourism and protected area facilities
- sustainable use of marine and coastal resources
- desertification and rangelands
- desertification and agro-biodiversity
- environmental legislation
- scientific research
- education and public awareness
- invasive species and GMOs
- environmental monitoring and impact assessments and
- multilateral environmental agreements.

The first and second strategic goals of the NBSAP deal with the development of management plans and the basic services to support such management plans in the protected areas. The first strategic goal is concerned with the development of protected areas, both terrestrial and marine. At this time, Qatar has five terrestrial protected areas and three marine protected areas. The terrestrial areas account for about 18% of the land surface of Qatar and include most of the habitat types. This strategic goal highlights the need for a protected area system plan and localized management plan as well as increasing public awareness of and participation in protection of the areas. The second strategic goal outlines the development of ecologically based tourism in the scenic areas of Qatar. This would include developing facilities and visitor centers at the sites. In addition local residents would act as guides and there would be an economic benefit to the local community with revenue sharing.

The third strategic goal is designed at protecting coastal and marine resources through the development of sustainable fishing and sustainable marine recreation. This would require the implementation of a coastal management plan as well as implementation of legislation regulating fishing and other activities that can threaten the marine resources.

Desertification is the focus of the fourth and fifth strategic goals. Given the seriousness of the threat of desertification in Qatar this seems appropriate. The fourth

strategic goal deals specifically with managing rangelands to prevent over-grazing, reducing the numbers of grazing animals and utilizing the traditional conservation ideas such as the *hima*.

The sixth strategic goal is to improve the environmental legislation and implement existing environmental legislation by improving the enforcement mechanisms.

The seventh and eighth strategic goals are aimed at increasing knowledge across all sectors of society from scholars to students. The seventh strategic goal aims to increase basic scientific research by increasing funding and supporting surveys with the goals of publishing the results. This will provide government leaders with the necessary information on which to base decisions that may significantly impact the environment. The eighth strategic goal focuses on increasing public awareness of biodiversity and improving environmental education through support of institutions and educators.

The ninth, tenth and eleventh strategic goals are concerned with utilizing the modern tools to control modern threats. The ninth strategic goal aims to protect the natural ecosystem, as well as humans, from the impacts of invasive species and genetically modified organisms (GMOs). This could be achieved by the establishment of a certification program for food and other imported products. The tenth goal is to strengthen the Environmental Impact Assessment process by finalizing rules, regulations and guidelines for the process. Finally, the eleventh strategic goal is to strengthen the multilateral and multinational environmental agreements by developing a planning process that would be comprehensive in fulfilling the requirements of these agreements. The dangers posed by GMOs and invasive species are of lower priority than the immediate threats to the marine and terrestrial environment (Third National Report, 2007).

The Third National Report on Implementation of the Convention on Biological Diversity (2007), essentially a report on implementation of the NBSAP, highlights a lack of trained experts as a limiting factor in implementing many areas of the Action Plan and Strategy. However, we can hope for increasing expertise since biological sciences/health and environmental studies do comprise 12% of MSc studies done abroad by Qataris. Biology, health, chemistry, and geology make up 24% of PhD studies being done abroad by Qataris (World Bank, 2005).

## Qatar's Protected Areas

In 2004 Qatar enacted the Law concerning the Protection of Wildlife and Their Natural Habitats. This law allows SCENR to propose protected areas but not designate protected areas. However, SCENR must protect species within the protected areas and can mandate regulations to do that. Unfortunately, guards deployed in protected areas are not trained in protected area management. The Protected Area Action Plan (2007) developed by SCENR outlines the current status of protected areas in Qatar as well as the actions to develop the Protected Areas management within the next five years (PAAP, 2007).

Qatar has five designated protected areas for the terrestrial ecosystems and three for the marine environment. These include the newly designated Al Reem biosphere reserve, Al Thakira, Shahanniya, Khor Al Udaid, Al Weseil, Al Oraiq and Al Mas'habiya (see Figure 2). Together they comprise about 18% of the terrestrial area of Qatar. Shahanniya, Al Mas'habiya and Al Isheiriq were wildlife farms developed into captive breeding sites for endangered species (Protected Area Action Plan (PAAP), 2007).

The maintenance and regulation of the Protected Areas of Qatar is expected to improve significantly.


Currently, a management plan is being developed for the Al Rheem biosphere, nearly 1200 square kilometres. Within the Al Reem area is about 18,000 people (PAAP, 2007). 75% of this population is males, mostly expatriate workers. Based on poverty statistics this would suggest that 40% of those expatriate workers survive under the poverty line. This may make it difficult to control illegal hunting and gathering activities. Most other protected areas, barring the captive breeding centers, have a resident population. Khor Al Udaid, at just over 1100 square kilometres and a unique geological site in the region, may soon be nominated as a UNESCO World Heritage Site. Unlike the other protected areas, there is no resident population within this protected area.

The Protected Area Action Plan (2007) outlines eleven steps that Qatar plans to implement for the successful development of the protected areas. This includes the following:

- Institutional reform for Protected Area (PA) management in order to streamline the management process
- Formation of a National Advisory Committee on PAs to coordinate agencies involved in the PAs
- Local consultative committees for PAs to address the concerns of the local population
- Categorize PAs as ex-situ conservation or in-situ conservation efforts to clarify management goals
- Preparation of a management plan for each PA
- Build capacity in protected area management
- Infrastructure development which aid the management of the PA
- Increase research activities such as species inventories, population monitoring, vegetation studies etc.
- Develop a sustainable resource use plan for each PA
- Institute a plan for the development of ecotourism
- Develop a policy guiding species reintroductions

Developments are expected to take place within SCENR which will contribute to improved conservation efforts. At this point, within SCENR, there is no Protected Areas Department. The Wildlife Management Department, within SCENR, is responsible for the administration of the protected areas. However, in the future expect there to be a department dealing strictly with Protected Areas and implementing the detailed management plan of each of the areas (SCENR, pers. Comm.).

Figure 2. Protected Areas in Qatar



© CIRS, Georgetown University School of Foreign Service in Qatar, 2008.

## Traditional Activities

The State of Qatar has placed an emphasis on preserving the environment so as to protect the health and livelihood of local communities (Bedouins) and looks toward traditional knowledge and culture when instituting national development plans (Third National Report, 2007). Part of this is maintaining viable land where traditional activities can be pursued. Traditional activities, as typically perceived, include fishing, agriculture and hunting. However, when we consider the economic aspect of traditional activities, Qataris are simply not involved in fishing, agriculture or hunting activities for economic benefit. It is the non-Qatari population that is taking part in the economic aspect of traditional practices (World Bank, 2005).

Traditional activities such as hunting the desert hare with Siluki dogs and falconry require both the land and animal populations. As development continues limiting access to areas that were once used for traditional activities, there is increasing conflict between Qataris wanting to engage in traditional activities and other individuals, activities or developments. Restoration and maintenance of the rangeland ecosystem, which could be accomplished in a relatively short time given favourable weather conditions, would provide the space and species for such activities to continue (Breulmann et. al., 2007). However, such measures would need to be instituted in a timely manner given the current rate of development. Winter camping is another traditional activity enjoyed by the local population and protected by the government. Appropriate management of winter camping and provision of facilities in areas of high use may be necessary to protect the fragile coastal environment and the species that inhabit that environment.

## Sustainable Development

Sustainable development, as first thought of by the 1987 Bruntland Commission report, describes sustainable development as development that “meets the needs of the present generation without compromising the ability of future generations to meet their own needs.” This can be thought of as economic and social development that does not threaten ecosystem services. However, in order to develop sustainably, a country must have some way to “measure” their development.

Qatar is currently in the process of developing a manageable system of sustainable development indicators for the country including social, economic and environmental indicators (Planning Council, 2006). These indicators are based on data collected from 2000-2005 and show clear trends in social, economic and environmental areas. Social indicators include a measure of social services that impact the standard of living of the population such as the provision of health care services and education. Economic indicators include aspects of basic economic structure such as macroeconomic performance as well as measure of gender equality and consumption (waste production). Environmental indicators include measurement of air quality, land degradation, water quantity and quality and biodiversity. The trends witnessed in these indicators are varied. For example, over the time period 2000-2005 fertilizer use remained about the same and pesticide use decreased slightly. However, the fish catch has almost doubled while protected areas have increased dramatically.

International sustainable development indicators are used in many countries and by international organizations to help guide sustainable development practices. The 2005 Environmental Stability Index allows for cross-national comparison of environmental progress and is a measure of overall progress toward environmental sustainability for the 142 countries involved (ESI, 2005). The scores are based on a core of 20 indicators, which may include 2-8 variables, for a total of 68 possible variables. However, Qatar has only 37 data points of the required minimum of 45. Of the total 238 countries initially analysed, Qatar was one of only 18 countries that fell into this data deficient category.

The development of sustainable development indicators is an important step in the development of Qatar. These indicators allow decision-makers to measure the progress of the country in regards to what they view as the most important factors in the development of the country. Stating clear goals to be attained would provide incentives for individuals, businesses, and decision-makers.



# What Limits Conservation Efforts in Qatar?

Conservation programs, in order to be successful, must be based on sound scientific knowledge such as distribution, occurrence, ecological function and importance (local, regional and international). This includes knowing what species are present, what their population numbers are and how they are changing, what the threats are to the species, and what the habitat requirements are for the species. Many well-intentioned conservation efforts have failed due to a lack of understanding of the species needs and the threats to the species. There must also be a willingness to participate in conservation measures at the cost of some activities.

The status and trends of endangered plants and animals is monitored by the IUCN, World Conservation Union. The IUCN publishes the Red List outlining the species, the status of the species, (near threatened, threatened, vulnerable, endangered, and critically endangered) and the population trend. As of 2004, Qatar had 33 species listed on the Red List. However, this is likely to be a gross underestimate because data are simply lacking for Qatar. The distribution, occurrence and local importance of many species are unknown. If the species listed from Kuwait, Bahrain, and the United Arab Emirates, which are likely to occur in Qatar, were included then the number of Red List species would increase to 67. Some of the species not listed for Qatar but are listed for neighbouring countries are certainly present in Qatar. One example is the whale shark (*Rhincodon typus*). In addition to this, those species that are listed are often data deficient, status unknown, or data out of date. For example, the critically endangered hawksbill turtle, for Qatar, population information is “out of date,” for the vulnerable dugong population information is “unknown,” for the finless porpoise there is “no data” ([www.iucnredlist.org](http://www.iucnredlist.org)). These are just a few of many examples. Finally, there are likely to be many species that have yet to be discovered. If an amateur group of naturalists can discover on a field trip a new species known to science (Macdonald, 2007), imagine what might be discovered if there were comprehensive, professional surveys conducted.

The scientific process ultimately depends on a strong foundation of knowledge, to which scientists continually contribute. Science builds on previous scientific studies; it is a cumulative process. Over the last 30 years there has been research on the fauna of Arabia but only a small portion of animals present have been officially recorded. The lack of basic research necessary for sustainable use and conservation programs is simply lacking. Conservation and sustainable use and development programs ultimately depend on a comprehensive scientific understanding of the species, its habitat requirements and the habitat itself (Brook et. al. 2006). Here in Qatar, there is a dearth of information about the natural environment. This can be confirmed by a search of the scientific literature concerning Qatar. A search on a common science search engine ([www.sciencedirect.com](http://www.sciencedirect.com)) returned 215 article hits for Qatar (263 for Bahrain, 557 for the UAE, 791 for Oman, and 1,123 for Kuwait). This may be one indication of the lack of scientific research concerning Qatar.

If Qatar is pursuing a science based decision-making process, then the amount of scientific work directly addressing Qatar will need to increase. Many of the studies that have been done have been performed by industry or individual research centers. The final product, the reports which contain the precious data, has not reached the internationally accessible scientific literature. As a result, new studies must start from the

very basics in order to build the public foundational knowledge necessary for conservation. At the least, publicizing hard to access studies and reports that have been done and making them openly available to scientists, using a resource such as the internet, would be a tremendous step forward for conservation in Qatar. This would require collaborative work between universities, government bodies, NGOs and industries. Such an effort, lead by government, combined with multi-year comprehensive studies to document the species present, their distribution and abundance would be another step towards a scientifically sounds and successful conservation strategy in Qatar.

An accessible base of ecosystem knowledge about Qatar, available in a format for decision-makers can significantly improve conservation efforts by the leadership. As stated earlier, in order to properly manage and protect the environment, a thorough understanding of the conditions and needs of the species is necessary. However, this information must be in a format that is usable by decision-makers. Recent publications such as the *Millennium Ecosystem Assessment* (MEA 2005) and the *Intergovernmental Panel on Climate Change* (IPCC, 2007) were designed specifically with the lawmaker in mind in order to facilitate the decision-making process. Without a well designed guideline for decision makers, it is important that development activities occur with the advice and cooperation of local research centers (Yasseen and Al-Thani, 2007). Research centers have been largely underutilized in their capacity to help make informed choices concerning planning as development proceeds.

## Conclusion

Conservation of species and maintaining a healthy ecosystem has global as well as local benefits. The State of Qatar plays an important role in the protection of some of the world's most critically endangered species, and supports many uniquely adapted organisms, some of which may still be unknown to science. However, these species and the habitats that support them are at risk from current development. Coastal development is particularly damaging to marine species and may threaten commercial fishing. This, combined with other factors such as desertification and soil salination due to water over-exploitation, demands a response. While the State of Qatar has moved quickly to ratify international conventions and establish the regulatory and managerial bodies, the implementation has been slow. This is in part due to a lack of trained workforce and scientific expertise available in the country. Yet, development of industrial and residential areas continues unabated at an increasing pace. Therefore, the development of an effective conservation program that is strictly implemented lags behind industrial and economic development. Much of the environmentally related work has been supported or lead by industry. A concerted effort lead by the government, combining financial and scientific support and a multi-year comprehensive terrestrial and marine survey, could greatly strengthen conservation efforts. Only then will we know what Qatar has to conserve, before it is too late.

# References

- Abderrahman, W.A. and Husain, T. (2006) "Pollution Impacts of Desalination on Ecosystems in the Arabian Peninsula." In *Policy Perspectives for Ecosystem and Water Management in the Arabian Peninsula*. Eds. Amer, K.M. Boer, B., Brook, M.C., Adeel, Z. Clusener-Godt, M. and Saleh, W. Hamilton, Canada: United Nations University International Network on Water, Environment and Health and UNESCO.
- Ahmed, I.F. (2002) *Qatar & the Sea*. Third edition. Doha, Qatar. National Council for Culture, Arts & Heritage.
- Al Habshi, A., Youssef, T., Aizpuru, M. and Basco, F. (2007) New mangrove Ecosystems Data Along the UAE Coast Using Remote Sensing. *Aquatic Ecosystem Health and Management* 10(3): 309-319.
- Al Yousef, A. H., Abdalla, O.A.S., and Akbar, M. (2000) *Progress Report for the National Report of the State of Qatar on the UNCCD Implementation*. State of Qatar Ministry of Municipal Affairs & Agriculture, Dept. of Agricultural & Water research Soil Research Section.
- Al-Ghadban, A.N. and Price, A.R.G. (2002) *Dredging and Infilling*. In *The Gulf Ecosystem: Health and Sustainability*. Eds. Khan, N.Y., Munawar, M. and Price, A.R.G. Backhuys.
- Barth, H-J. (2007) *Crab Induced Salt Marsh regeneration after the 1991 Gulf War Oil Spill*. *Aquatic Ecosystem Health and Management* 10(3): 309-319.
- Bourn, D. (2003) *Livestock Dynamics in the Arabian Peninsula: A Regional Review of National Livestock Resources and International Livestock Trade*. Environmental Research Group Oxford.
- Breulmann, M., Boer, B. Wernery, U., Wernery, R. El Shaer, H., Alhadrami, G., Gllacher, D., Peacock, J., Ali Chaudhary, A., Brown, G. and Norton, J. (2007) *The Camel: From Tradition to Modern Times*. A Proposal Towards Combating Desertification via the Establishment of Camel Farms Based on Fodder Production from Indigenous Plants and Halophytes. UNESCO.
- Brook, M.C., Al Shoukri, S., Amer, K.M., Boer, B. and Krupp, F. (2006) "Physical and Environmental Setting of the Arabian Peninsula and Surrounding Seas." In *Policy Perspectives for Ecosystem and Water Management in the Arabian Peninsula*. Eds. Amer, K.M. Boer, B., Brook, M.C., Adeel, Z. Clusener-Godt, M. and Saleh, W.

Brook, M.C., Al Houqani, H. and Al Mugrin, A. (2006) "The Current Status and Future Requirements of Water Resources Management in the Arabian Peninsula." In *Policy Perspectives for Ecosystem and Water Management in the Arabian Peninsula*. Eds. Amer, K.M. Boer, B., Brook, M.C., Adeel, Z. Clusener-Godt, M. and Saleh, W. Hamilton, Canada: United Nations University International Network on Water, Environment and Health and UNESCO.

Carpenter, K.E., Krupp, F., Jones, D.A. and Zajonz, U. (1997) *Living Marine Resources of Kuwait, Eastern Saudi Arabia, Bahrain, Qatar and the United Arab Emirates*. Rome, FAO.

Child, S. (2006) "Biosalinity Research and Development in the Arabian Peninsula Countries." In *Policy Perspectives for Ecosystem and Water Management in the Arabian Peninsula*. Eds. Amer, K.M. Boer, B., Brook, M.C., Adeel, Z. Clusener-Godt, M. and Saleh, W. Hamilton, Canada: United Nations University International Network on Water, Environment and Health and UNESCO.

Borwn, G. Peacock, J. Loughland, R. and Alhadrami, G. (2006) "Coastal and Terrestrial Ecosystem Management Requirements in the Arabian Peninsula." In *Policy Perspectives for Ecosystem and Water Management in the Arabian Peninsula*. Eds. Amer, K.M. Boer, B., Brook, M.C., Adeel, Z. Clusener-Godt, M. and Saleh, W. Hamilton, Canada: United Nations University International Network on Water, Environment and Health and UNESCO.

Chaudhary, J. R., and Husain, T. (2006) "Uncertainty Analysis of Humidity and Precipitation Changes Using Data from Global Climactic Models with a Case Study." *EIC Climate Change Technology*, 2006 IEEE, May 2006, 1-10.

Dietz, S. and Adger, W.N. (2003) "Economic Growth, Biodiversity Loss and Conservation Effort." *Journal of Environmental Management* 68:23-35.

El-Sayed Selim, M. (2004) *Environmental Security in the Arab World*. Meeting of the International Studies Association, 17-20 March 2004, Montreal, Canada.

FAO (2003) *Information on Fisheries Management in the State of Qatar*. <http://www.fao.org/fi/fcp/en/QAT/body.htm>.

Gillespi, F. (2007) "Qatar Natural History Group Discover New Marine Species!" *Gulf Times*, 15 September 2007.

Gulf Times (2008) "Qatar's Population Tops 1.5 million." *Gulf Times*, 29 January 2008.

Jones, D.A., Ealey, T., Baca, B., Livesey, S. and Al-Jamali, F. (2007) "Gulf Desert Developments Encompassing a Marine Environment, a Compensatory Solution to the Loss of Coastal Habitats by Infill and Reclamation: The case of the Pearl City Al-Khiran, Kuwait." *Aquatic Ecosystem Health and Management* 10(3):268-276.

- Khan, N.Y. (2007) "Multiple Stressors and Ecosystem-based Management in the Gulf." *Aquatic Ecosystem Health and Management* 10(3): 259-267.
- Khan, N.Y. (2002) "Physical and Human Geography." In *The Gulf Ecosystem: Health and Sustainability*. Eds. Khan, N.Y., Munawar, M. and Price, A.R.G. Backhuys.
- Krupp, F., Al-Muftah, A., Jones, D.A. and Hoolihan, J. (2006) "Marine and Coastal Ecosystem management Requirements in the Arabian Peninsula with special regard to water resources." In *Policy Perspectives for Ecosystem and Water Management in the Arabian Peninsula*. Eds. Amer, K.M. Boer, B., Brook, M.C., Adeel, Z. Clusener-Godt, M. and Saleh, W. Hamilton, Canada: United Nations University International Network on Water, Environment and Health and UNESCO.
- Loughland, R.A., Darwish, A.M., Saddiqui, K., Fadel, S.S., Al-Ali, A.A. and Crawford, K. (2007) *The Development of a Marine and Coastal Natural Resource Atlas: Abu Dhabi Emirate, United Arab Emirates*.
- Mitchell, T. and Hulme, M. (2000) *A Country-by-Country Analysis of Past and Future Warming Rates*. Tyndall Centre Working Paper No. 1, November 2000, UEA, Norwich, UK, 6pp.
- National Biodiversity Strategy and Action Plan* (2004) State of Qatar [www.cbd.int/countries/?country=qa](http://www.cbd.int/countries/?country=qa).
- The Peninsula* (2007) "Steady Growth in Annual Fish Catch." 11 July 2007.
- Planning Council (2007) *Human Development Report for the State of Qatar*. The Planning Council – General Secretariat.
- Planning Council (2006) *Sustainable Development Indicators in Qatar*. The Planning Council – General Secretariat.
- Protected Area Management Plan 2008-2013* (2007) State of Qatar. Department of Wildlife Conservation, SCENR.
- Rezai, H., Wilson, S., Claerebolid, M., and Riegl, B. (2004) "Coral Reef Status in the ROPME Sea Area: Arabian/Persian Gulf, Gulf of Oman and Arabian Sea" in *Status of the Coral Reefs of the World*.
- Riegl, B. (2002) "Effects of the 1996 and 1998 positive Sea Surface Temperature Anomalies on Corals, Coral Diseases and Fish in the Arabian Gulf" (Dubai, UAE) *Marine Biology* 140: 29-40.
- Seddon, P. and Mallon, D. (2007) Arabian Conservation Workshops. *Oryx* 41(2) 131-134.
- Sheppard, C. and Loughland, R. (2002) "Coral Mortality and Recovery in response to increasing temperature in the southern Arabian Gulf." *Aquatic Ecosystems and Health Management* 5(4): 395-402.

Sheppard, C., Price, A. and Roberts, C. (1992) *Marine Ecology of the Arabian Peninsula: Patterns and processes in Extreme Tropical Environments*. Academic Press, Harcourt Brace Jovanovich, San Diego, CA.

Sheppard, C. (2003) "Predicted recurrences of mass coral mortality in the Indian Ocean." *Nature* 425: 294-297.

SOMER (2003) *State of the Marine Environment Report*. ROPME/GC-11/003. Regional Organization for the Protection of the Marine Environment, Kuwait, 217 pp.

Third National Report (2007) "On Implementation of the Convention on Biological Diversity" [www.cbd.int/countries/?country=qa](http://www.cbd.int/countries/?country=qa).

World Bank (2005) *Labor market Strategy for the State of Qatar: Main Report*. Volume 1 Dec 29, 2005.

Yasseen, B.T., and Al-Thani, R.F. (2007) "Halophytes and associated properties of natural soils in the Doha area, Qatar." *Aquatic Ecosystem Health & Management* 10:3, 320-326.

Zhang, X., Aguilar, E., Sensoy, S., Melkonyan, H., Tagoyeva, U., Ahmed, N., Kutaladze, N., Rhimzadeh, F., Taghipour, A., Hantosh, T., Albert, P., Semawi, M., Karam, Ali, M., Halal Said Al-Shabibi, M., Al-Oulan, Z., Zatar, T., Al Dean Khelet, I., Hamoud, S., Sagar, R., Demircan, M., Eken, M., Adiguzel, M., Alexander, L., Peterson, T., and Wallis, T. (2005) "Trends in Middle East climate extreme indices from 1950-2003." *Journal of Geophysical Research*, Vol. 110, D22104, doi:10.1029/2005JD006181, 2005.

Center for International and Regional Studies

[sfs.georgetown.edu/qatar/about/cirs](https://sfs.georgetown.edu/qatar/about/cirs)







CIRS

CENTER FOR  
INTERNATIONAL  
AND REGIONAL  
STUDIES